


Forschungsschwerpunkt S92

# Industrial Geometry

<http://www.industrial-geometry.at>

---



FSP Report No. 91

**Volumes with Piecewise Quadratic  
Medial Surface Transforms:  
Computation of Boundaries and  
Trimmed Offsets**

B. Bastl, B. Jüttler, J. Kosinka, M. Lávička

November 2009

---

**FWF**

Der Wissenschaftsfonds.



# Volumes with Piecewise Quadratic Medial Surface Transforms: Computation of Boundaries and Trimmed Offsets

Bohumír Bastl<sup>a</sup>, Bert Jüttler<sup>b,1</sup>, Jiří Kosinka<sup>c</sup> and Miroslav Lávička<sup>a</sup>

<sup>a</sup> University of West Bohemia, Faculty of Applied Sciences, Department of Mathematics,  
Univerzitní 8, 301 00 Plzeň, Czech Republic

<sup>b</sup> Johannes Kepler University, Institute of Applied Geometry,  
Altenberger Str. 69, 4040 Linz, Austria

<sup>c</sup> University of Oslo, Centre of Mathematics for Applications,  
P.O. Box 1053 Blindern, NO-0316 Oslo, Norway

## Abstract

MOS surfaces are rational surfaces in  $\mathbb{R}^{3,1}$  which provide rational envelopes of the associated two-parameter family of spheres. Moreover, all the offsets admit rational parameterizations as well. Recently, it has been proved that quadratic triangular Bézier patches in  $\mathbb{R}^{3,1}$  are MOS surfaces. Following this result, we describe an algorithm for computing an exact rational envelope of a two-parameter family of spheres given by a quadratic patch in  $\mathbb{R}^{3,1}$ . The main focus of this paper is given to geometric aspects of the algorithm. Since these patches are capable of producing  $C^1$  smooth approximations of medial surface transforms of spatial domains, we use this algorithm to generate rational approximations of envelopes of general medial surface transforms. One of the main advantages of this approach to offsetting is the fact that the trimming procedure becomes considerably simpler.

Keywords: Quadratic Bézier triangles, MOS surfaces, trimmed offsets

## 1 Introduction

Generating valid tool paths in NURBS form has become a universal standard in technical applications such as CNC machining in recent years. Hence, shape (curve, surface or volume) offsets and corresponding algorithms have been widely studied in Computer-Aided Design and Manufacturing, see [47, 27, 15, 36, 48, 53, 54, 55] and references therein for more details. However, free-form NURBS shapes do not possess rational offsets in general and thus suitable approximation techniques, often based on rational curves or surfaces of relatively low degree, are used – see e.g. [17, 14, 32, 59, 61, 57]. Nevertheless, these offset approximation techniques suffer from several shortcomings. First, one needs to approximate each offset independently. Moreover, the constructed approximation of offsets to the same base shape may not have a constant distance to each other, which is not acceptable in some applications, e.g. architecture. Second, although a rational approximation of an offset is computed, the so called trimming, i.e., the detection and elimination of self-intersections,

is still a very difficult problem. Let us emphasize that the approximation can possess self-intersections even though the original offset has none. Even in the case of curves this leads to challenging computational and time-consuming problems and the situation is even more complicated for surfaces, cf. [58, 56, 40]. The purpose of this paper is to identify an approach that avoids these problems.

The approximate techniques for offsets are now widely used in CAD systems since they are capable of dealing with the problems appearing in technical practice mentioned above – but usually at the expense of great computational effort. Therefore, it is worthwhile to investigate exact techniques as well, i.e., to study shapes with exact rational offsets. These can be subsequently used for formulating suitable approximation techniques for free-form shapes. Compared to the classical approximation techniques, not offsets but the base shape is approximated and it is guaranteed that all corresponding offsets are rational. Therefore, only one approximation step is required even if more than one offset is needed. Moreover, all the offsets are at a constant distance from each other.

Shapes with rational offsets have been studied for many years. In the case of planar curves, the class of Pythagorean Hodograph (PH) curves as polynomial curves possessing rational offset curves and polynomial arc-length functions was introduced in [21]. A thorough analysis of PH curves has followed – see e.g. [20, 25, 18, 59, Pelosi et al.(2007)]. Later, the concept of polynomial planar PH curves was generalized to space PH curves ([22, Farouki et al.(2003), Farouki and Sakkalis(2007), Šír and Jüttler(2007)]) and to rational PH curves ([49, 50, 43]). Analogously, the notion of rational surfaces with rational offsets, the so called Pythagorean Normal vector (PN) surfaces, was introduced in [49]. More details about PN surfaces can be found in [35, 31, 33]. For a survey of shapes with Pythagorean normals property (i.e., possessing rational offsets) see [19].

Later, it has been proved in [Sampoli et al.(2006)] that surfaces with Linear field of Normal vectors (LN surfaces), introduced in [Jüttler(1998)], provide rational convolution surfaces with an arbitrary rational surface. Since spheres admit rational descriptions, LN surfaces possess exact rational offsets. This

<sup>1</sup>Corresponding author. E-mail: bert.juettler@jku.at, phone/fax: +43 732 2468 9178 / 29162, Homepage: www.ag.jku.at.

result was applied in [Jüttler and Sampoli(2000)] to constructing exact rational offsets of LN surfaces. As observed recently ([Lávička and Bastl(2007), 44]), all non-developable polynomial quadratic surfaces belong to the class of LN surfaces and therefore possess rational offsets. Motivated by the fact that approximation by quadratic Bézier surfaces provides a good compromise between curvature reproduction and computational cost and offers an extra feature, i.e., the PN property, these patches were used in [2] for formulating an algorithm for generating rational approximations to offsets of general free-form surfaces.

However, even though that PH curves and PN surfaces admit rational offsets, the usually most costly ingredient, the trimming, still needs to be performed as in the case of approximation techniques. An alternative approach to the problem in the curve case based on the medial axis transform (MAT) of a planar domain, introduced in [4], was formulated in [37] and [10]. There is a one-to-one correspondence between the MAT in three-dimensional Minkowski space  $\mathbb{R}^{2,1}$  and the object boundary, i.e., for a given geometric object there is a unique MAT and conversely, the object can be directly reconstructed from its MAT. The main advantages of this description follow from the dimensional reduction while topological properties are maintained. As observed in [10, 37], if a segment of the medial axis transform is an MPH (Minkowski Pythagorean Hodograph) curve, then the associated branches of the domain boundaries are segments of rational curves. In addition, all offsets of the domain boundaries have also this property. A thorough analysis of MPH curves has followed in [8, 9, 11, 26, 28, 30]. Moreover, using the MAT representation makes the trimming procedure for the inner offsets very simple – only those parts of the MAT where the corresponding circle radius is less than the offset distance  $\delta$  have to be trimmed. For more details see e.g. [34, 52, 12, 13, 7, 6, 1].

The situation in three-dimensional space has become an active research area recently, since the so called MOS surfaces as a spatial analogy of MPH curves were introduced in [29]. The medial surface transform (MST) of a volume is the set of surface patches (or curve segments) in four-dimensional Minkowski space  $\mathbb{R}^{3,1}$  such that each point of these surfaces/curves represents the center and the radius of a maximal sphere inscribed into the domain. The MST covers the structure of the domain and thus can be used in all sorts of geometric modelers (constructive solid geometry (CSG) and boundary representation (B-rep)) as a useful tool. The distinguishing property of MOS surfaces is that if considered as an MST of a volume, the associated envelope and its offsets admit exact rational parameterization. Later, it was proved in [42] that quadratic triangular Bézier surfaces in  $\mathbb{R}^{3,1}$  possess the MOS property and a related study followed in [41, 45].

Following the results of [42], [2] and [3], we describe an efficient algorithm for computing boundaries and trimmed offsets of volumes with piecewise quadratic medial surface transforms. Considering general (free-form) medial surface transforms in  $\mathbb{R}^{3,1}$ , we produce their  $C^1$  approximations using quadratic Bézier patches (cf. [39, 60]) which gives us an extra feature – the MOS property. Thus we have a guarantee on the rationality of the volume boundaries as well as all offsets. The presented technique also simplifies the computation of self-intersections of inner offsets and the follow-up trimming procedure considerably.

The remainder of this paper is organized as follows. Section

2 recalls some basic facts concerning medial surface transforms, envelopes of two-parameter family of spheres and MOS surfaces. Section 3 is devoted to quadratic triangular Bézier surfaces in  $\mathbb{R}^{3,1}$ , their (isotropic) normals and isotropic Gauss images. In this section, we also formulate and discuss an algorithm for envelope computation for quadratic MSTs. The algorithm is then demonstrated on several examples in Section 4. Finally, we conclude the paper.

## 2 Preliminaries

We recall the relation between the four-dimensional Minkowski space  $\mathbb{R}^{3,1}$  and the medial surface transform (MST) of a volume. We then consider rational patches on sheets of the MST and the computation of the corresponding patches on the boundary surface of the volume.

### 2.1 Medial surface transform

The *four-dimensional Minkowski space*  $\mathbb{R}^{3,1}$  is the four-dimensional real affine space which is equipped with the indefinite inner product

$$\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u}^\top \mathbf{J} \mathbf{v} = u_1 v_1 + u_2 v_2 + u_3 v_3 - u_4 v_4 \quad (1)$$

where  $\mathbf{J} = \text{diag}(1, 1, 1, -1)$ . The hyperplane  $x_4 = 0$  is a three-dimensional subspace, and the restriction of the inner product to this hyperplane is the usual Euclidean inner product. We identify this hyperplane with the *three-dimensional Euclidean space*. More precisely, any point  $\mathbf{x} = (x_1, x_2, x_3)^\top \in \mathbb{R}^3$  is equivalently considered as a point  $\mathbf{x} = (x_1, x_2, x_3, 0)^\top \in \mathbb{R}^{3,1}$ .

For any point  $\mathbf{p} = (p_1, p_2, p_3, p_4)^\top \in \mathbb{R}^{3,1}$ , let  $\mathbf{p}' = (p_1, p_2, p_3)^\top$  be its projection into  $\mathbb{R}^3$ . The set

$$\mathcal{B}_{\mathbf{p}} = \{\mathbf{x} \in \mathbb{R}^3 : \langle \mathbf{p} - \mathbf{x}, \mathbf{p} - \mathbf{x} \rangle \leq 0\} \quad (2)$$

is the *oriented ball* with center  $\mathbf{p}'$  and radius  $|p_4|$ . The sign of the radius  $p_4$  defines the orientation of the ball. Its boundary is the *oriented sphere*  $\mathcal{S}_{\mathbf{p}} = \delta \mathcal{B}_{\mathbf{p}}$  which contains all points  $\mathbf{x} \in \mathbb{R}^3$  satisfying

$$\langle \mathbf{p} - \mathbf{x}, \mathbf{p} - \mathbf{x} \rangle = 0. \quad (3)$$

Any oriented ball in  $\mathbb{R}^3$  can be identified with a unique point in the four-dimensional Minkowski space  $\mathbb{R}^{3,1}$ .

We consider a bounded volume  $V$  with a smooth boundary  $\delta V$ . The set of all balls with non-negative radius (i.e., with non-negative orientation) which are contained in  $V$  is partially ordered with respect to inclusion. The maximal elements of this set form the *medial surface transform* (MST) of  $V$ , cf. Fig. 1. We consider them as points in Minkowski space, which then form the set  $\text{MST}(V) \subset \mathbb{R}^{3,1}$ .

The medial surface transform of the volume  $V$  consists of components of dimensions two, one, and zero, which are called *sheets*, *seams* and *junctions*, respectively. Their points correspond to maximal inscribed balls which – in the generic case – touch the boundary in two, three, and four points, respectively. The sheets meet in seams, and the seams meet in junctions.

In the remainder of this paper we consider a two-dimensional rational surface patch

$$\mathbf{p} : \Omega \rightarrow \mathbb{R}^{3,1} : (u, v) \mapsto \mathbf{p}(u, v) \quad (4)$$




Figure 1: Bisector surface (blue) and two branches (red and green) of the envelope surface forming a boundary of a volume  $V$ .

with domain  $\Omega \subseteq \mathbb{R}^2$  and parameters  $u, v$ , which defines a rational parameterization of a subset of the interior of a sheet. Every ball  $\mathcal{B}_{\mathbf{p}(u,v)}$  touches the boundary  $\delta V$  of the volume in two points. The contact points form two surface patches on the boundary. The centers  $\mathbf{p}'(u, v)$  of the balls form the *bisector surface* of these two surface patches – see Fig. 1.

## 2.2 Envelopes of two-parameter families of spheres

Given a patch (4) of the medial surface transform, we are interested in computing the associated patches of the boundary surface, see Fig. 1. These two patches are obtained as the *envelope* of the two-parameter family of spheres

$$\{\mathcal{S}_{\mathbf{p}(u,v)} : (u, v) \in \Omega\}. \quad (5)$$

By differentiating the equation (3) of the sphere we obtain two equations

$$\langle \partial_u \mathbf{p}, \mathbf{p} - \mathbf{x} \rangle = 0, \quad \langle \partial_v \mathbf{p}, \mathbf{p} - \mathbf{x} \rangle = 0 \quad (6)$$

which – along with the sphere equation (3) – characterize the points  $\mathbf{x}$  of the envelope surfaces. Recall that  $\mathbf{x} \in \mathbb{R}^3$  is a point in the hyperplane  $x_4 = 0$ . The substitution

$$\mathbf{x} = \mathbf{p} - \frac{p_4}{n_4} \mathbf{n} \quad (7)$$

transforms the three equations (3) and (6) into a homogeneous non-linear system of equations

$$\langle \mathbf{n}, \mathbf{n} \rangle = 0 \quad \text{and} \quad (8)$$

$$\langle \mathbf{n}, \partial_u \mathbf{p} \rangle = 0, \quad \langle \mathbf{n}, \partial_v \mathbf{p} \rangle = 0 \quad (9)$$

for the vector  $\mathbf{n} \in \mathbb{R}^{3,1}$ .

The two equations (9) express the fact that the vector  $\mathbf{n}$  is *orthogonal* to the two-dimensional tangent plane of the surface patch  $\mathbf{p}$  with respect to the Minkowski inner product (1). The first equation (8) restricts the solutions to *isotropic* vectors (vectors of zero length with respect to the Minkowski metric). Consequently, the non-trivial solutions of the system are the *isotropic normal vectors* of the surface.

Summing up, the points  $\mathbf{x}$  of the envelope surfaces can be generated by

1. finding all non-trivial isotropic normal vectors and
2. computing  $\mathbf{x}$  from (7).

Note that any nontrivial solution of (8) satisfies  $n_4 \neq 0$ .

## 2.3 MOS surfaces

The sub-determinants<sup>2</sup>

$$P_{ij} = \det \begin{pmatrix} \partial_u p_i & \partial_v p_i \\ \partial_u p_j & \partial_v p_j \end{pmatrix}, \quad i, j = 1, \dots, 4, \quad (10)$$

of the two  $4 \times 2$  matrix  $(\partial_u \mathbf{p}, \partial_v \mathbf{p})$  formed by the partial derivative vectors can be used to express the solutions of the system (8)–(9),

$$\mathbf{n}_{\pm}(\mu) = \mu \begin{pmatrix} P_{12}P_{24} + P_{13}P_{34} \pm \sqrt{C}P_{23} \\ P_{21}P_{14} + P_{23}P_{34} \pm \sqrt{C}P_{31} \\ P_{31}P_{14} + P_{32}P_{24} \pm \sqrt{C}P_{12} \\ -(P_{23}^2 + P_{31}^2 + P_{12}^2) \end{pmatrix}, \quad \mu \in \mathbb{R}, \quad (11)$$

with

$$C = P_{12}^2 + P_{31}^2 + P_{23}^2 - P_{14}^2 - P_{24}^2 - P_{34}^2. \quad (12)$$

The number of linearly independent solutions depends on the sign of  $C$ , as follows.

- If  $C > 0$ , then we obtain two families of solutions. The orthogonal complement space of the two-dimensional tangent plane, which is described by (9), intersects the quadratic cone of isotropic vectors (8) in two lines. We obtain two points of the envelope surfaces. The tangent plane is said to be *space-like*.
- If  $C = 0$ , then we obtain one family of solutions. The orthogonal complement space of the two-dimensional tangent plane intersects the quadratic cone of isotropic vectors (8) in one line. We obtain one point of the envelope surfaces. The tangent plane is said to be *light-like*.
- If  $C < 0$ , then we obtain no non-trivial solutions. The orthogonal complement space of the two-dimensional tangent plane intersects the quadratic cone of isotropic vectors only in the origin, which is the apex of the cone. The tangent plane is said to be *time-like*.

If the surface patch (4) describes a subset of the interior of a MST sheet, then  $C(u, v) > 0$  is satisfied for all  $(u, v) \in \Omega$ . All points in the domain correspond to space-like tangent planes.

<sup>2</sup>These sub-determinants are the Plücker coordinate of the line (cf. [51]) in the plane at infinity, which is spanned by the two derivative vectors.

Consequently, for all  $(u, v) \in \Omega$ , two solutions (11) with two associated points of the envelope surface exist. Moreover, the bisector surface is regular, since

$$\|\partial_u \mathbf{p}' \times \partial_v \mathbf{p}'\|^2 = P_{23}^2 + P_{31}^2 + P_{12}^2 \geq C. \quad (13)$$

Finally we recall a definition from [29] which generalizes the notion of Minkowski Pythagorean hodograph (MPH) curves. If there exists a polynomial  $\sigma(u, v)$  such that  $C = \sigma^2$ , then the parameterization of the envelope surfaces, which is obtained by combining (11) with (7), is rational. The given surface  $\mathbf{p}$  is then a *MOS surface*, i.e., a *Medial surface Obeying the Sum of squares condition*

$$C = \sigma^2 \iff P_{12}^2 + P_{31}^2 + P_{23}^2 = \sigma^2 + P_{14}^2 + P_{24}^2 + P_{34}^2. \quad (14)$$

### 3 Quadratic patches as medial surface transforms

A quadratic patch in  $\mathbb{R}^{3,1}$  is defined by the power basis representation

$$\mathbf{p}(u, v) = \frac{1}{2} \mathbf{q}_{20} u^2 + \mathbf{q}_{11} uv + \frac{1}{2} \mathbf{q}_{02} v^2 + \mathbf{q}_{10} u + \mathbf{q}_{01} v + \mathbf{q}_{00}, \quad (15)$$

with the coefficient vectors  $\mathbf{q}_{ij} \in \mathbb{R}^4$ , where the parameters  $u, v$  span a suitable domain. We will mostly work with the standard triangular domain  $\Delta \subset \mathbb{R}^2$  given by  $u \in [0, 1]$  and  $v \in [0, 1 - u]$ . According to the affine classification of quadratic patches [46], there exist 42 different types of quadratic patches. In the remainder of this paper we exclude developable patches (planes, cones and parabolic cylinders) from our considerations.

#### 3.1 Normals

The two equations (9) take the form

$$\begin{aligned} f_{\mathbf{n}}(u, v) &= \langle \mathbf{n}, \mathbf{q}_{20} \rangle u + \langle \mathbf{n}, \mathbf{q}_{11} \rangle v + \langle \mathbf{n}, \mathbf{q}_{10} \rangle = 0, \\ g_{\mathbf{n}}(u, v) &= \langle \mathbf{n}, \mathbf{q}_{11} \rangle u + \langle \mathbf{n}, \mathbf{q}_{02} \rangle v + \langle \mathbf{n}, \mathbf{q}_{01} \rangle = 0. \end{aligned} \quad (16)$$

For any given normal  $\mathbf{n}$ , they form a linear system for the parameters  $(u, v)$ .

The number of solutions  $(u, v)$  depends on the normal  $\mathbf{n}$ . Since the system is homogeneous with respect to  $\mathbf{n}$ , we consider the normals as points in a three-dimensional real projective space

$$\mathcal{N} = P^3(\mathbb{R}). \quad (17)$$

This three-dimensional space is the hyperplane at infinity of the four-dimensional Minkowski space.

Let

$$d(\mathbf{n}) = \mathbf{n}^\top \mathbf{J}(\mathbf{q}_{20} \mathbf{q}_{02}^\top - \mathbf{q}_{11} \mathbf{q}_{11}^\top) \mathbf{J} \mathbf{n} \quad (18)$$

be the determinant of the linear system (16). Further, let

$$\begin{aligned} d_u(\mathbf{n}) &= \mathbf{n}^\top \mathbf{J}(\mathbf{q}_{11} \mathbf{q}_{01}^\top - \mathbf{q}_{02} \mathbf{q}_{10}^\top) \mathbf{J} \mathbf{n}, \\ d_v(\mathbf{n}) &= \mathbf{n}^\top \mathbf{J}(\mathbf{q}_{10} \mathbf{q}_{11}^\top - \mathbf{q}_{01} \mathbf{q}_{20}^\top) \mathbf{J} \mathbf{n} \end{aligned} \quad (19)$$

be the two determinants generated by applying Cramer's rule.

First we describe the relation between the normals  $\mathbf{n}$  and the number of solutions of the system (16).

- The set

$$\mathcal{S} = \{\mathbf{n} : d(\mathbf{n}) = 0\}, \quad (20)$$

is called the *set of singular normals*. It is a singular quadric surface in  $\mathcal{N}$ .  $\mathcal{S}$  consists of all normals with no or non-unique solutions of (16). Consequently, if  $\mathbf{n} \in \mathcal{N} \setminus \mathcal{S}$ , then the set of solutions consists of a single point in the  $uv$ -plane.

- The set

$$\mathcal{E} = \{\mathbf{n} : d(\mathbf{n}) = d_u(\mathbf{n}) = d_v(\mathbf{n}) = 0\}, \quad (21)$$

which is contained in  $\mathcal{S}$ , is called the *set of exceptional normals*. It is a (possibly degenerate) cubic space curve, which is defined as the intersection of three quadric surfaces.  $\mathcal{E}$  consists of all normals where several solutions of (16) exist. If  $\mathbf{n} \in \mathcal{E}$ , then the set of all solutions forms a line in the  $uv$ -plane.

#### 3.2 The fibration of $\mathcal{N}$

Next we study the sets

$$\mathcal{F}_{(u,v)} = \{\mathbf{n} \in \mathcal{N} : f_{\mathbf{n}}(u, v) = g_{\mathbf{n}}(u, v) = 0\} \quad (22)$$

of normals which satisfy the two equations (16) for given values of  $(u, v)$ . They will be called the *fibers*. They form a partition of  $\mathcal{N} \setminus \mathcal{S}$  into mutually disjoint subsets.

- If the quadratic surface is regular at  $(u, v)$ , then the fiber  $\mathcal{F}_{(u,v)}$  is a line in  $\mathcal{N}$ . Indeed, it is defined as the intersection of the two linearly independent planes (9). At singular points, the fiber is a plane (if  $\text{rank}(\partial_u \mathbf{p}, \partial_v \mathbf{p}) = 1$ ) or even the entire space  $\mathcal{N}$  (if both derivatives vanish). The latter case occurs only for conical quadratic surfaces, which were excluded.
- Any line which is contained in a fiber  $\mathcal{F}_{(u,v)}$  intersects the set  $\mathcal{E}$  of exceptional normals in two (possibly conjugate-complex) points (counted with multiplicities). Indeed, the two intersections of the line with the singular set  $\mathcal{S}$  satisfy the equations for the given values  $(u, v)$ , hence these intersections belong to the exceptional set, too.

**Example 1** We discuss several examples which are taken from the affine classification of quadratic patches in [46].

1. Let us consider a non-hyperplanar patch given by

$$\mathbf{p}_1(u, v) = \left( \left( u + \frac{3}{2} \right)^2 + v, v^2, \left( u + \frac{3}{2} \right) v, u + \frac{3}{2} \right)^\top.$$

The exceptional set of this patch is a cubic space curve (see Fig. 2, top row).

2. Let us consider another non-hyperplanar patch

$$\mathbf{p}_2(u, v) = \left( -u^2 + u \left( v + \frac{1}{2} \right) + v^2 + v + \frac{1}{4}, u^2 + u + v^2 + 2v + \frac{3}{4}, u^2 + u - v^2 + \frac{1}{4}, -v - \frac{1}{2} \right)^\top.$$

The exceptional set is a conic and a line in this case (see Fig. 2, middle row).




Figure 2: Left: Exceptional lines (thin blue), the curve  $C(u, v) = 0$  (thick blue) and the domain triangle  $\Delta$ . Right: The curve  $\mathcal{S}_\Sigma$  (green), exceptional isotropic normals (green points) and the exceptional set  $\mathcal{E}$  (blue). Top row: Example 1, patch  $\mathbf{p}_1(u, v)$ ; Middle row: Example 1, patch  $\mathbf{p}_2(u, v)$ ; Bottom row: Example 1, patch  $\mathbf{p}_3(u, v)$ .

3. Let us consider a hyperplanar patch

$$\mathbf{p}_3(u, v) = \begin{pmatrix} \frac{1}{400} (289u^2 + 17u(17v + 40) + 289v^2 + 85v + 75, \\ -289u^2 - 289uv + 289v^2 + 595v + 325, \\ 289u^2 + 17u(17v + 20) - 5(51v + 25), \\ (17u - 15)(17u - 17v - 5) \end{pmatrix}^\top,$$

The exceptional set consists of three lines (see Fig. 2, bottom row).

### 3.3 Isotropic normals

So far we studied solutions of the system of equations (9) for a quadratic patch (15). Now we will analyze the consequences of the additional equation (8) for isotropic normals.

The set of isotropic normals (i.e., the set of normals satisfying (8)) forms an oval quadric  $\Sigma$  in the three-dimensional projective space  $\mathcal{N}$ . Without loss of generality we may assume that the isotropic normals satisfy  $n_4 = 1$ . The oval quadric of isotropic normals can then be identified with the unit sphere

$$0 = \langle \mathbf{n}, \mathbf{n} \rangle = n_1^2 + n_2^2 + n_3^2 - 1. \quad (23)$$

We analyze the relation between the isotropic normals and the number of solutions of the system (8)–(9).

- The intersection curve of the set  $\mathcal{S}$  and the isotropic quadric  $\Sigma$  is the spherical quartic curve  $\mathcal{S}_\Sigma = \mathcal{S} \cap \Sigma$  of *isotropic singular normals*, cf. Fig. 3 (right). It consists of all isotropic normals with no or non-unique solutions of (9). Consequently, if  $\mathbf{n} \in \Sigma \setminus \mathcal{S}_\Sigma$ , then the set of solutions consists of a single point in the  $uv$ -plane.
- The intersection points of the sphere of isotropic normals with the exceptional curve  $\mathcal{E}$ ,  $\mathcal{E}_\Sigma = \mathcal{E} \cap \Sigma$  are the – at most 6 – *exceptional isotropic normals*, Fig. 3 (right). If  $\mathbf{n} \in \mathcal{E}$ , then the set of all solutions forms a line in the  $uv$ -plane. We obtain at most 6 lines in the  $uv$ -plane, which will be called the *exceptional lines* in the parameter domain – see Fig. 3 (left).
- Consider a fiber  $\mathcal{F}_{(u,v)}$  at a regular point  $\mathbf{p}(u, v)$ . If  $C(u, v)$  is positive, zero, or negative, then the fiber intersect the unit sphere in two, one, or no points. The curve defined by  $C(u, v) = 0$  in the parameter domain thus corresponds to a spherical curve  $\mathcal{L}_\Sigma$  where the fibers touch  $\Sigma$ . This is the *curve of isotropic normals at points with light-like tangent planes* – see Fig. 3 (right).

### 3.4 The isotropic Gauss images

Let us assume that the domain  $\Omega$  of the surface patch contains only points satisfying  $C(u, v) > 0$ . Consequently, all tangent planes are space-like, and all points are regular. This assumption is satisfied for surfaces which represent a patch of the interior of a sheet of the MST.

For each point  $(u, v)$  we obtain two isotropic normals satisfying  $n_4 = 1$ , by choosing  $\mu = -1/(P_{23}^2 + P_{31}^2 + P_{12}^2)$  in Eq. (11). They are exactly the two intersections of the fiber  $\mathcal{F}_{(u,v)}$  with  $\Sigma$ . Thus, the isotropic normals define a one-to-two mapping

$$\Omega \rightarrow \Sigma : (u, v) \mapsto \{\mathbf{n}_+(u, v), \mathbf{n}_-(u, v)\} = \mathcal{F}_{(u,v)} \cap \Sigma. \quad (24)$$

The set of all isotropic normals defines the *isotropic Gauss image*  $\Gamma = \mathbf{n}_\pm(\Omega)$  of the given quadratic patch with domain  $\Omega$ .

If we consider the two isotropic normals along an exceptional line in the parameter domain, then we have the following two possibilities.

- Case 1: One isotropic normal is constant along the line, while the second isotropic normal varies. The constant isotropic normal is an exceptional one with the property that several fibers  $\mathcal{F}_{(u,v)}$  intersect in it.
- Case 2: Both isotropic normals are constant along the line. All points  $(u, v)$  of the exceptional line possess the same fiber  $\mathcal{F}_{(u,v)}$ , and this fiber intersects the sphere  $\Sigma$  in two exceptional isotropic normals. Consequently, the fiber is contained in  $\mathcal{E}$ .

Finally, let  $\Gamma_0 \subset \Sigma$  be the isotropic Gauss image without isotropic normals along the exceptional lines, and let  $\Omega_0$  be the parameter domain without exceptional lines. The isotropic normals map each connected component of  $\Omega_0$  to two connected components of  $\Gamma_0$  – see Fig. 3. When restricted to one of the two isotropic normals, this mapping is bijective.




Figure 3: Left: Exceptional lines (thin blue), the curve  $C(u, v) = 0$  (thick blue) and the domain triangle  $\Delta$ . Right: The two components of the isotropic Gauss image (red and blue), the curve  $\mathcal{S}_\Sigma$  (green), exceptional isotropic normals (green points) and the curve  $\mathcal{L}_\Sigma$  (black).

### 3.5 Envelope computation

We describe an algorithm for computing the exact rational envelope of a 2-parameter family of spheres given by a quadratic patch (15) over  $\Delta$ . Its input is a quadratic Bézier triangle (15), where all points are assumed to be space-like, i.e.,  $C(u, v) > 0$  holds for all  $(u, v) \in \Delta$ . The algorithm proceeds in four steps.

**Step 1: Subdivision along exceptional lines** Compute the exceptional lines and triangulate the domain such that the exceptional lines are edges of the triangulation. For each triangle of the triangulation, apply a linear re-parameterization of the quadratic patch such that the parameter domain is again the standard triangle  $\Delta$ . If the exceptional lines do not intersect the interior of the parameter domain, then no subdivision is required (see Fig. 4).




Figure 4: Left: exceptional lines (thin blue), the curve  $C(u, v) = 0$  (thick red) and the domain triangle  $\Delta$ . Right: The two components of the isotropic Gauss image (red and blue), the curve  $\mathcal{S}_\Sigma$  (green) and exceptional isotropic normals (green points).

This step is similar to Algorithm 1 in [2], but the role of the parabolic lines is played by the exceptional ones. The next three steps are applied to all quadratic patches which are generated by Step 1. For these patches, at most one of the three boundary curves is an exceptional line.

**Step 2: Covering the isotropic Gauss images** For each of the patches obtained in Step 1, compute the isotropic Gauss images.

These Gauss images are either two spherical triangles or biangles, or a biangle and a triangle, all with curved boundaries. The biangles occur if one of the domain boundaries is an exceptional line. Next we compute rational spherical patches

$$\mathbf{n}^* : \Delta \rightarrow \Sigma : (s, t) \mapsto \mathbf{n}^*(s, t) \quad (25)$$

which cover the Gauss images. In the case of a biangle, one may use singular patches where the singular point coincides with the image of the exceptional line. These rational spherical patches are found using stereographic projection (see Fig. 5).

This step is similar to Algorithm 2 in [2]. As the main difference we have to consider the two isotropic Gauss images instead of only one Gauss image.




Figure 5: Left: Stereographic projection of one component of the isotropic Gauss image (black) and circumscribed triangle (green). Right: One component of the isotropic Gauss image (red) and its covering patch (green).

**Step 3: Envelope computation** For each spherical rational patch  $\mathbf{n}^*$  we compute one segment of the envelope from (7), where  $\mathbf{n}$  is replaced with  $\mathbf{n}^*$  and the parameters  $(u, v)$  of the patch  $\mathbf{p}$  are replaced with

$$u(s, t) = \frac{d_u(\mathbf{n}^*(s, t))}{d(\mathbf{n}^*(s, t))}, \quad v(s, t) = \frac{d_v(\mathbf{n}^*(s, t))}{d(\mathbf{n}^*(s, t))}, \quad (26)$$

cf. (18) and (19). This gives rational triangular patches of degree 10. The surface patches obtained by applying the substitution (26) to (15) are MOS surfaces.

In the case of a biangular isotropic Gauss image, the use of a spherical patch with a singular point ensures that the envelope surface possesses a regular rational parameterization, which is obtained after omitting a common factor of numerator and denominator, similar to the Example presented in Section 4.3 of [2].

**Step 4: Trimming** The spherical patches  $\mathbf{n}^*$ , which cover the isotropic Gauss images, are generally larger than these images, i.e., they may contain isotropic normal vectors which do not correspond to isotropic normal vectors of the MST patch over  $\Delta$ . Hence, we need to restrict  $\Delta$  to an appropriate subset given by polynomial inequalities resulting from

$$u(s, t) \geq 0 \wedge v(s, t) \geq 0 \wedge 1 - u(s, t) - v(s, t) \geq 0, \quad (27)$$

which can be derived from (26), see Fig. 6.




Figure 6: Left: Trimmed parameter domain of the envelope patches bounded by brown, orange and yellow curves. Right: Bisector surface (blue) and two branches of the envelope patch (red and green).

Summing up, the envelope surface consists of the trimmed patches which are generated by applying the steps 2, 3 and 4 to all triangular patches obtained from Step 1 of this algorithm.

The computation of the exceptional lines leads to a polynomial equation of degree 6, hence the solutions cannot be found in closed form. Consequently, even if the coefficients of the input patch are rational, then the parameterizations of the envelopes which are generated by the algorithm do not possess rational coefficients.

Parameterizations with rational coefficients can be generated by using a simplified version of the algorithm, which omits the subdivision step and uses spherical patches covering the isotropic Gauss images which do not necessarily possess singular points at the exceptional isotropic normals. However, the resulting parameterizations of the envelope surfaces may possess singularities and points where both numerators denominator vanish simultaneously.

We have implemented the algorithm using floating point numbers. In order to avoid numerical problems, we used the Bernstein-Bézier representations of the triangular patches. Bounds on the error introduced by an approximate computation of the exceptional lines and several examples have been presented in [3].

## 4 MST approximation and trimming of offset surfaces

We discuss the approximation of patches on the sheets of the MST of a general volume by piecewise quadratic surfaces. This approximate representation of the MST is then used for the computation of trimmed offset surfaces.

More precisely, we consider a surface patch  $s : [0, 1]^2 \rightarrow \mathbb{R}^3$  which represents a part of the boundary of the domain  $V$  and which corresponds to a patch of a sheet of the MST. Consequently, for each point  $s(u, v)$ , there is a maximal inscribed ball which touches the boundary  $\delta V$  in this point and in another point. We use the approach described in [16] for numerically computing the centers  $c(u, v)$  and the radii  $r(u, v)$  of the maximal inscribed balls, i.e., for evaluating the points of the MST.

After generating a set of values (centers of the spheres and radii), we use it to construct an approximation of the MST by

a piecewise quadratic spline surface  $\mathbf{P} : [0, 1]^2 \rightarrow \mathbb{R}^{3,1}$ , which is defined over a criss-cross triangulation of the domain. See [5, 60, 39] for more information on this type of spline functions. Our implementation uses a global least-squares fitting of the sampled points. Alternatively, one may use quasi-interpolation methods, such as the techniques described in [23, 24]. If the grid is sufficiently fine, then the piecewise quadratic approximation is guaranteed to possess only space-like tangent planes, since the first derivatives of the MST are approximated, too.

We can now apply the exact envelope computation method, which was described in the previous section, to each polynomial segment  $\mathbf{p}$  of the quadratic spline surface  $\mathbf{P}$ . Recall that the trimmed inner offsets of the boundary surface of a domain can be obtained by (1) subtracting the offsetting distance from the 4th component of the MST (which represents the radius of the maximal inscribed balls) and (2) restricting the modified MST to points with non-negative fourth components. Consequently, if the piecewise quadratic approximation  $\mathbf{P}$  of the MST is available, we can use it for computing exact trimmed offsets of the boundary surface of the volume which is represented by  $\mathbf{P}$ . We implemented this by adding another trimming procedure to our algorithm, which restricts the piecewise quadratic MST to the domain where the radius is non-negative.

**Example 2** We consider the volume  $V \subset \mathbb{R}^3$  bounded by the two surface patches

$$\begin{aligned} \mathbf{s}_1(u, v) &= \left(u, v, \frac{1}{3}uv\right)^\top, \\ \mathbf{s}_2(u, v) &= \left(u, v, \frac{1}{3}(u^2 + v^2) + 1\right)^\top, \end{aligned} \quad (28)$$

$(u, v) \in [-1, 1]^2$ , and approximate the associated MST by a piecewise quadratic surface consisting of 12 triangular patches. The maximum distance error is equal to 3.28% of the diameter of the bounding box. For each of these patches we parameterize the envelope surfaces as described in the previous section. All 12 patches of the approximating quadratic spline surface possess only space-like points and moreover no additional subdivisions of the parameter domains are needed. Finally, Fig. 7 shows the two branches of the approximating envelope (red and green), the quadratic spline approximation of the medial surface (blue) and the two branches of inner offsets (light red and light green). In case of the self-intersecting inner offset, the trimmed regions on the envelope are illustrated.

**Example 3** We consider another volume in  $\mathbb{R}^3$ . Similarly to the previous example, we approximate its MST by a quadratic spline which consists of 24 triangular patches. The maximum distance error is equal to 1.83% of the diameter of the bounding box. Again, for each of these patches we parameterize the envelope. All patches possess only space-like points and no additional subdivision is needed. Fig. 8 shows the two branches of the approximating envelope (red and green), the quadratic spline approximation of the medial surface (blue) and two branches of inner offsets (yellow and purple) which are trimmed if the self-intersection occurs.




Figure 7: Bisector surface (blue) and the two branches of the envelope (red and green) in Example 2, along with the trimmed inner offsets (light red and light green) – without self-intersection (top right) and with self-intersection (bottom left and right).

## 5 Conclusion

In this paper we presented an algorithm for computing boundaries and trimmed offsets of volumes given by piecewise quadratic medial surface transforms. Since polynomial quadratic patches in  $\mathbb{R}^{3,1}$  belong to the class of MOS surfaces and are capable of producing  $C^1$  approximations to free-form surfaces considered as medial surface transforms, our algorithm can be used for computing rational approximations of volume boundaries and all their offsets. Only one approximation step is required, and the trimming procedure for inner offsets is greatly simplified using this approach.

**Acknowledgments** B. Bastl and M. Lávička were supported by Research Plan MSM 4977751301. B. Bastl, B. Jüttler and M. Lávička were supported by grant AKTION 2009/5 (MEB060905).

## References

- [1] Aichholzer, O., Aigner, W., Aurenhammer, F., Hackl, T., Jüttler, B., Rabl, M.: *Medial axis computation for planar free-form shapes*. Computer-Aided Design, Volume 41, Issue 5, Pages 339–349, 2009.
- [2] B. Bastl, B. Jüttler, J. Kosinka, and M. Lávička: *Computing exact rational offsets of quadratic triangular Bézier surface patches*. Computer-Aided Design, Vol. 40, pp. 197–209, Elsevier, 2008.
- [3] Bastl, B., Kosinka, J., Lávička, M.: *A Symbolic-Numerical Envelope Algorithm Using Quadratic MOS Patches*. Proceedings of SIAM/ACM Joint Conference on Geometric and Physical Modeling, 2009.
- [4] H. Blum: *A transformation for extracting new descriptors of shape*. In: W. Wathen-Dunn (Ed.), Models for the perception of speech and visual form, pp. 362–380, MIT Press, 1967.
- [5] de Boor, C., Höllig, K., Riemenschneider, S.: *Box Splines*. Springer, 1993.
- [6] L. Cao, Z. Jia, and J. Liu: *Computation of medial axis and offset curves of curved boundaries in planar domains based on the Cesaro’s approach*. Computer Aided Geometric Design, Volume 26, Issue 4, May 2009, Pages 444–454, 2009.
- [7] L. Cao and J. Liu: *Computation of medial axis and offset curves of curved boundaries in planar domain*. Computer-Aided Design, Vol. 40, pp. 465–475, Elsevier, 2008.
- [8] H.Ch. Cho, H.I. Choi, S.-H. Kwon, D.S. Lee, and N.-S. Wee: *Clifford algebra, Lorentzian geometry and rational parametrization of*




Figure 8: Bisector surface (blue) and the two branches of the envelope (red and green) in Example 3, along with the trimmed inner offsets (yellow and purple) – without self-intersection (top right) and with self-intersection (bottom left and right).

- canal surfaces. *Computer Aided Geometric Design*, Vol. 21, pp. 327–339, Elsevier, 2004.
- [9] H.I. Choi, S.W. Choi, and H.P. Moon: *Mathematical theory of medial axis transform*. *Pacific Journal of Mathematics*, Vol. 181, pp. 57–88, 1997.
- [10] Choi, H. I., Han, Ch. Y., Moon, H. P., Roh, K. H., Wee, N. S.: *Medial axis transform and offset curves by Minkowski Pythagorean hodograph curves*. *Computer-Aided Design* **31**, 1999, 59–72.
- [11] H.I. Choi, D.S. Lee, and H.P. Moon: *Clifford algebra, spin representation and rational parameterization of curves and surfaces*. *Advances in Computational Mathematics*, Vol. 17, pp. 5–48, 2002.
- [12] Culver, T., Keyser, J., Manocha, D.: *Exact computation of the medial axis of a polyhedron*. *Comput. Aided Geom. Des.* **21** (2004), 65–98.
- [13] W.L.F. Degen: *Exploiting curvatures to compute the medial axis for domains with smooth boundary*. *Computer Aided Geometric Design*, Vol. 21, pp. 641–660, Elsevier, 2004.
- [14] Elber G, Lee I-K, Kim M-S. Comparing Offset Curve Approximation Methods. *IEEE Comp Graphics and Appl* 1998; 17: 62–71.
- [15] Elber G, Kim M-S. Special issue on Offsets, sweeps, and Minkowski sums. *Computer-Aided Design* 1999; 31: 163.
- [16] Elber, G., and Kim, M. S.: *A Computational Model for Non-rational Bisector Surfaces: Curve-Surface and Surface-Surface Bisectors*. *Geometric Modeling and Processing* 2000, Hong Kong, pp 364–372.
- [17] Farouki RT. The approximation of non-degenerate offset surfaces, *Computer Aided Geometric Design* 1986; 3:15-43.
- [18] Farouki, R. T., *Pythagorean-hodograph curves*. *Handbook of Computer Aided Geometric Design* (J. Hoschek, G. Farin and M.-S. Kim, eds.), Elsevier, 2002, 405–427.
- [19] R.T.: Farouki, *Pythagorean-Hodograph Curves: Algebra and Geometry Inseparable*. *Geometry and Computing*. Springer-Verlag, 2008.
- [20] Farouki, R.T., and Neff, C.A., (1995), *Hermite interpolation by Pythagorean-hodograph quintics*. *Math. Comput.* 64, 1589–1609.
- [21] R.T. Farouki, T. Sakkalis: *Pythagorean hodographs*. *IBM Journal of Research and Development*, Vol. 34, pp. 736–752, 1990.
- [22] Farouki, R.T. and Sakkalis, T., *Pythagorean-hodograph space curves*. *Adv. Comput. Math.* 2 (1994) 41–66.
- [Farouki and Sakkalis(2007)] Farouki, R.T., Sakkalis, T.: *Rational space curves are not “unit speed”*. *Computer Aided Geometric Design*, Vol. 24, pp. 238–240. Elsevier, 2007.
- [Farouki et al.(2003)] Farouki, R.T., Manni, C. and Sestini, A.: *Spatial  $C^2$  PH quintic splines*. In: Lynche, T., Mazure, M.L. and Schumaker, L.L. (eds.), *Curve and Surface Design: St. Malo 2002*, pp. 147–156. Nashboro Press, 2003.
- [23] Foucher, F. and Sablonnière, P.: *Approximating partial derivatives of first and second order by quadratic spline quasi-interpolants on uniform meshes*. *Mathematics and Computers in Simulation* **77** (2008), 202–208.
- [24] Foucher, F. and Sablonnière, P.: *Quadratic spline quasinext term-interpolants and collocation methods*. *Mathematics and Computers in Simulation*, Volume 79, Issue 12, Pages 3455–3465, 2009
- [Jüttler(1998)] Jüttler, B.: *Triangular Bézier surface patches with linear normal vector field*. In: Cripps, R. (ed.), *The Mathematics of Surfaces VIII*, pp. 431–446. Information Geometers, 1998.
- [Jüttler and Sampoli(2000)] Jüttler, B., Sampoli, M.L.: *Hermite interpolation by piecewise polynomial surfaces with rational offsets*. *Computer Aided Geometric Design*, Vol. 17, pp. 361–385. Elsevier, 2000.
- [25] Jüttler, B. (2001), *Hermite interpolation by Pythagorean hodograph curves of degree seven*. *Math. Comp.* 70, 1089–1111.
- [26] G.-I. Kim and M.-H. Ahn:  *$C^1$  Hermite interpolation using MPH quartic*. *Computer Aided Geometric Design*, Vol. 20, pp. 469–492, 2003.
- [27] Kimmel R, Bruckstein AM. Shape offsets via level sets, *Computer-Aided Design* 1993; 25: 154-62.

- [28] J. Kosinka and B. Jüttler:  *$G^1$  Hermite Interpolation by Minkowski Pythagorean hodograph cubics*. Computer Aided Geometric Design, Vol. 23, pp. 401–418, Elsevier, 2006.
- [29] Kosinka, J., Jüttler, B.: *MOS Surfaces: Medial surface transforms with rational domain boundaries*. Lecture Notes in Computer Science, Vol. 4647, pp. 245–262. Springer, 2007.
- [30] J. Kosinka and B. Jüttler:  *$C^1$  Hermite interpolation by Pythagorean hodograph quintics in Minkowski space*. Advances in Computational Mathematics, Vol. 30, pp. 123–140, 2009.
- [31] R. Krasauskas: *Branching blend of natural quadrics based on surfaces with rational offsets*. Computer Aided Geometric Design, Vol. 25, pp. 332–341, Elsevier, 2008.
- [32] Kulczycka MA, Nachman LJ. Qualitative and quantitative comparisons of B-spline offset surface approximation methods, Computer-Aided Design 2002; 34: 19-26.
- [Lávička and Bastl(2007)] Lávička, M., Bastl, B.: *Rational hypersurfaces with rational convolutions*. Computer Aided Geometric Design, Vol. 24, pp. 410–426. Elsevier, 2007.
- [33] M. Lávička and B. Bastl: *PN surfaces and their convolutions with rational surfaces*. Computer Aided Geometric Design, Vol. 25, pp. 763–774, Elsevier, 2008.
- [34] D.T. Lee: *Medial axis transformation of a planar shape*. IEEE Trans Pattern Analysis and Machine Intelligence, Vol. PAMI-4, pp. 362–369, 1982.
- [35] W. Lü and H. Pottmann: *Rational parameterization of quadrics and their offsets*. Computing, Vol. 57, pp. 135–147. Springer, 1996.
- [36] Maekawa T. An overview of offset curves and surfaces. Computer-Aided Design 1999; 31: 165–73.
- [37] Moon, H. P.: *Minkowski Pythagorean hodographs*. Computer Aided Geometric Design **16**, (1999), 739–753.
- [38] T. Nishita, T. W. Sederberg, and M. Kakimoto: *Ray Tracing Trimmed Rational Surface Patches*. ACM SIGGRAPH Computer Graphics, Vol. 24, pp. 337–345, 1990.
- [39] Nürnberger G., Zeilfelder F.,: *Developments in bivariate spline interpolation*. Journal of Computational and Applied Mathematics 2000; 121: 125-52.
- [40] Diana Pekerman, Gershon Elber, Myung-Soo Kim: *Self-intersection detection and elimination in freeform curves and surfaces*. Computer-Aided Design, Volume 40, Issue 2, Pages 150-159, 2008.
- [Pelosi et al.(2007)] Pelosi, F., Sampoli, M.L., Farouki, R.T. and Manni, C.: *A control polygon scheme for design of planar  $C^2$  PH quintic spline curves*. Computer Aided Geometric Design, Vol. 24, pp. 28–52. Elsevier, 2007.
- [41] M. Peternell and B. Odehnal: *On Generalized LN-Surfaces in 4-Space*. In: Proceedings of 'ISSAC08', pp. 223–230, 2008.
- [42] Peternell, M., Odehnal, B. and Sampoli, M.L.: *On quadratic two-parameter families of spheres and their envelopes*. Computer Aided Geometric Design, Vol. 25, pp. 342–355. Elsevier, 2008.
- [43] M. Peternell and H. Pottmann: *A Laguerre Geometric Approach to Rational Offsets*. Computer Aided Geometric Design, Vol. 15, pp. 223–249, Elsevier, 1998.
- [44] Peternell M, Odehnal B.: *Convolution surfaces of quadratic triangular Bézier surfaces*. Computer Aided Geometric Design, Vol. 25, pp. 116–129. Elsevier, 2008.
- [45] Peternell, M.: *Rational two-parameter families of spheres and rational offset surfaces*. Journal of Symbolic Computation, In Press.
- [46] Peters, J., Reif, U.: *The 42 equivalence classes of quadratic surfaces in affine n-space*. Computer Aided Geometric Design 1998, 15: 459-73.
- [47] Pham B. Offset curves and surfaces: a brief survey. Computer-Aided Design 1992; 24: 223-9.
- [48] Piegl LA, Tiller W. Computing offsets of NURBS curves and surfaces. Computer-Aided Design 1999; 31: 147-156.
- [49] H. Pottmann: *Rational curves and surfaces with rational offsets*. Computer Aided Geometric Design, Vol. 12, pp. 175–192, Elsevier, 1995.
- [50] H. Pottmann and M. Peternell: *Applications of Laguerre geometry in CAGD*. Computer Aided Geometric Design, Vol. 15, pp. 165–186, Elsevier, 1998.
- [51] Pottmann, H., Wallner, J.: *Computational Line Geometry*. Springer, Berlin 2001.
- [52] M. Ramanathan and B. Gurumoorthy: *Constructing medial axis transform of planar domains with curved boundaries*. Computer-Aided Design, Vol. 35, pp. 619–632, Elsevier, 2003.
- [53] Ravi Kumar GVV, Shastry KG, Prakash BG. Computing non-self-intersecting offsets of NURBS surfaces. Computer-Aided Design 2002; 34: 209-28.
- [54] Ravi Kumar GVV, Shastry KG, Prakash BG. Computing offsets of trimmed NURBS surfaces, Computer-Aided Design 2003; 35: 411-20.
- [55] Ravi Kumar GVV, Shastry KG, Prakash BG. Computing constant offsets of a NURBS B-Rep. Computer-Aided Design 2003; 35: 935–944.
- [Sampoli et al.(2006)] Sampoli, M.L., Peternell, M., Jüttler, B.: *Rational surfaces with linear normals and their convolutions with rational surfaces*. Computer Aided Geometric Design, Vol. 23, pp. 179–192. Elsevier, 2006.
- [56] Seong J-K, Elber G, Kim M-S. Trimming local and global self-intersections in offset curves/surfaces using distance maps, Computer-Aided Design 2006; 38: 183-93.
- [57] J.-L. Shih, S.-H. Frank Chuang: *One-sided offset approximation of freeform curves for interference-free NURBS machining* Computer-Aided Design, Volume 40, Issue 9, Pages 931–937.
- [58] Sun VF, Nee AYC, Lee KS. Modifying free-formed NURBS curves and surfaces for offsetting without local self-intersection. Computer-Aided Design 2004; 36: 1161-9.
- [59] Šír Z, Feichtinger R, Jüttler B. Approximating curves and their offsets using biarcs and Pythagorean hodograph quintics, Computer-Aided Design 2006; 38: 608-18.
- [Šír and Jüttler(2007)] Šír, Z. and Jüttler, B.:  *$C^2$  Hermite interpolation by Pythagorean hodograph space curves*. Mathematics of Computation, Vol. 76, pp. 1373-1391, 2007.
- [60] Wang, R. H., *Multivariate spline functions and their applications*. Dordrecht: Kluwer, 2001.
- [61] Zhao H-Y, Wang G-J. Error analysis of reparametrization based approaches for curve offsetting. Computer-Aided Design 2007; 39: 142-8.